

**NON-CONTRASTED COMPUTED TOMOGRAPHY FOR THE
ACCURATE MEASUREMENT OF LIVER STEATOSIS IN
NON ALCOHOLIC FATTY LIVER DISEASE**

Dissertation submitted for

**D.M.DEGREE EXAMINATION- AUGUST-2013
BRANCH-IV-MEDICAL GASTROENTEROLOGY
MADRAS MEDICAL COLLEGE**

&

**RAJIV GANDHI GOVERNMENT GENERAL HOSPITAL
CHENNAI-600003**



**THE TAMILNADU DR.M.G.R MEDICAL UNIVERSITY
CHENNAI-600032**

CERTIFICATE

This is to certify that the dissertation **“NON-CONTRASTED COMPUTED TOMOGRAPHY FOR THE ACCURATE MEASUREMENT OF LIVER STEATOSIS IN NON ALCOHOLIC FATTY LIVER DISEASE”** is a bonafide work of **Dr.P.SENTHIL KUMAR** in partial fulfillment of the requirements for D.M. Branch-IV (MEDICAL GASTROENTEROLOGY) examination of THE TAMILNADU DR.M.G.R. MEDICAL UNIVERSITY to be held in August 2013. The period of post-graduate study and training was from August 2010 to July 2013.

DR. MOHAMMED ALLI, MD., DM.,
PROFESSOR AND HOD
MADRAS MEDICAL COLLEGE,
CHENNAI

DEAN
MADRAS MEDICAL COLLEGE,
CHENNAI

DECLARATION

I, Dr. P.SENTHIL KUMAR, solemnly declare that this dissertation entitled, **NON-CONTRASTED COMPUTED TOMOGRAPHY FOR THE ACCURATE MEASUREMENT OF LIVER STEATOSIS IN NON ALCOHOLIC FATTY LIVER DISEASE** is a bonafide work done by me at the Department of Medical Gastroenterology, Madras Medical College & Rajiv Gandhi Government General Hospital, Chennai during the period 2010- 2013 under the guidance and supervision of the Prof. Mohammed Ali M.D.,D.M., Professor and Head of the Department Of Medical Gastroenterology of Madras Medical College & Government General Hospital, Chennai.

This dissertation is submitted to The Tamil Nadu Dr.M.G.R Medical University, towards requirement for the award of D.M. Degree (Branch-IV) in Medical Gastroenterology

Chennai
Date

Dr. P.SENTHIL KUMAR
Post graduate student
D.M(Medical Gastroenterology)
Madras Medical college, Chennai-3.

ACKNOWLEDGEMENTS

A great many people made this work possible. I thank my Dean for allowing me to conduct this study. I owe my warmest respects and sincere thanks to our beloved Professor Prof. Mohammed Ali Professor and Head of the Department of Medical Gastroenterology, Government General Hospital, Chennai, and Dr P. Padmanaban M.D., D.M., T. Pugazhenti M.D., D.M., additional professor Department of Medical Gastroenterology, Government General Hospitals Chennai who were the driving force behind this study. I am indebted to my Assistant Professors Dr. K. Prem Kumar, Dr. Kani Sheik Mohammed, Dr. Ratnakar Kini P, and who helped in my work

In addition, I am grateful to of my copost graduate students for helping me out throughout this study period. Last but not the least I thank all my patients for their kind cooperation

CONTENTS

S.NO	TITLE	PAGE NO
1	INTRODUCTION	1
2	AIM OF THE STUDY	6
3	REVIEW OF LITERATURE	7
4	MATERIALS AND METHODS	34
5	RESULTS	40
6	DISCUSSION	54
7	CONCLUSION	59
BIBLIOGRAPHY		
ANNEXURES		
➤ ABBREVIATIONS		
➤ CONSENT FORM		
➤ MASTER CHART		
➤ ETHICAL COMMITTEE APPROVAL ORDER		
➤ TURNITIN-PLAGIARISM SCREEN SHOT		
➤ DIGITAL RECEIPT		

ABBRIVATIONS

ncCT	-	Non-Contrasted Computed Tomography
LFT	-	Liver Function Test
CTHU	-	Computer Tomography in Houns Field Unit.
CTL-S	-	Computer Tomography of Liver – spleen in Houns Field Unit.
CTL/S	-	Computer Tomography of Liver to spleen ratio in Houns Field Unit.
BMI	-	Body Mass Index
NASH	-	Non Alcoholic Steato Hepatitis
NAFLD	-	Non Alcoholic Fatty Liver Disease

INTRODUCTION

INTRODUCTION

Non alcoholic fatty liver disease[NAFLD] is emerging as most common liver disorder in India and other developing countries, The histological spectrum ranging from simple steatosis or steatosis with only mild inflammation (type 1 and 2 NAFLD) to more severe steatohepatitis (types 3 and 4 NAFLD or NASH). Types 3 and 4 NAFLD progress to cirrhosis in 15-20% of patients. Progression is silent or paradoxically associated with normalization of aminotransferases. “NASH” coined by Ludwig and colleagues from Mayo clinic.

SPECTRUM OF NAFLD

- Steatosis
- Steatohepatitis (NASH)
- NASH with Fibrosis
- Cirrhosis

DEFINITIONS

NAFLD: Non alcoholic fatty infiltration with no or minimal inflammation and no fibrosis.also called simple steatosis,Pure steatosis and bland steatosis.

Primary NAFLD : indicates typical NAFLD associated with central obesity and T2 DM without specific etiology.

Secondary NAFLD – associated with a specific, non alcohol related problem such as drugs or toxins.

Toxin- associated steatohepatitis : associated with specific toxin or medication.eg petrochemical exposure in oil industry

CRITERIA FOR DEFINING NAFLD

- 1) A liver biopsy showing moderate to gross macrovesicular fatty change with or without inflammation (lobular or portal),and Mallory bodies, fibrosis, or cirrhosis.
- 2) No alcohol consumption or consumption less than 40 g of ethanol per week.
- 3) Absence of serologic evidence of hepatitis B or hepatitis C.

EPIDEMIOLOGY AND PREVALENCE

Prevalence of NAFLD 13-18% and that of NASH specifically 2-3% (1.2-9%) in two Japanese studies- incidence rate of 31 and 86 cases . (AGA guidelines 2012)

The prevalence of NAFLD defined by USG -46%

By histology confirmed NASH-12.2 %. (AGA guidelines 2012) It is a disease of all sexes, ethnicities, and age groups (peak 40-59) Higher in Hispanics than Africa Americans. Occurs more frequently in females (65 to 83%)

OBESITY

4.6 fold increase risk of fatty liver in USG compared to nonobese. Autopsy studies, steatosis-approx 70%obese vs 35% lean. Severe obesity patients- prevalence of NAFLD may go upto 90%. Obese patients with abnormal liver enzymes approx 30% has septal fibrosis and 10% cirrhosis.

INSULIN RESISTANCE AND T2 DM

Insulin resistance is very common in NAFLD. Progression to overt diabetes is preceded by steatosis in susceptible population. 75% of T2 DM- fatty infiltration. Liver injury worsens the preexistent diabetes in patients with NAFLD and doubles the prevalence of cirrhosis from 10-25%.

HYPERLIPIDEMIA

2/3 of hypertriglyceridemia&1/3rd with hypercholesterolemia- fatty liver Hyperlipidemia reported in 92% of NASH patients. Ethnic variation in lipid metabolism. Heterogenicity in lipid phenotypes in NAFLD

METABOLIC SYNDROME

Steatosis and central adiposity with underlying insulin resistance and lipotoxicity- independent risk factor for metabolic syndrome. Unexplained elevation of liver enzymes attributable to NAFLD seen in 7% of individuals with metabolic syndrome defined by adult Treatment Panel 111 criteria.

DIAGNOSIS

On routine liver biochemical test elevated liver enzymes level usually will suspect common causes of liver disease will arise for other investigations like USG, CT, MRI, CBC, PT, anti-HCV, HBsAg, serum iron, anti-trypsin, ANA, on physical examination hepatomegaly, on history alcohol consumption should be excluded [less than 40 grams /week] clinically if the patients having age > 50 years, obesity, diabetes, hypertension

IMAGING

Hepatic ultrasonogram shows “bright liver” with increased echoes consistent with steatosis on CT scan. Fatty liver shows lower in density compared to normal, MRI shows fat appears bright on T1 – weighted image, combination USG and CT have sensitivity 93 – 100% for detecting

hepatic fat, with more than grade I fatty liver [$< 5\%$ is minimal, $5 - 33\%$ grade I, $33 - 66\%$ grade II, $> 66\%$ grade III] with positive predictive value of $62 - 76\%$, no radiologic modality able to distinguish simple steatosis from more advanced NAFLD. These imaging techniques [USG, CT, MRI, support the diagnosis of NAFLD but it cannot predict the severity of disease. And also it cannot replace gold standard liver biopsy and histopathology, for establishing diagnosis in certainty.

So The degree of steatosis in NAFLD patients is usually assessed with invasive technique like liver biopsy, because of the invasive method of the liver biopsy and its painful nature of the procedure with moderate complication rate need to find alternative less invasive, less painful, quick method to accurately diagnose NAFLD

AIM OF THE STUDY

AIMS AND OBJECTIVES

To study and analyse unenhanced CT [nc-CT] determination of hepatic steatosis based on image attenuation data (Hounsfield units) and to correlate it with liver histopathology

REVIEW OF LITERATURE

REVIEW OF LITERATURE

NAFLD is emerging as one of the most common liver disorder in India and many other developing countries in the world. The histological spectrum ranging from simple steatosis or steatosis with only mild inflammation(type 1 and 2 NAFLD) to more severe steatohepatitis (types 3 and 4 NAFLD or NASH. Types 3 and 4 NAFLD progress to cirrhosis in 15-20% of patients. Progression is silent or paradoxically associated with normalization of aminotransferases. “NASH” coined by Ludwig and colleagues from Mayo clinic.

SPECTRUM OF NAFLD

- Steatosis
- Steatohepatitis (NASH)
- NASH with Fibrosis
- Cirrhosis

DEFINITE

NAFLD: Non alcoholic fatty infiltration with no or minimal inflammation and no fibrosis. Also called simple steatosis, pure steatosis and bland steatosis.

Primary NAFLD: indicates typical NAFLD associated with central obesity and T2 DM without specific etiology.

Secondary NAFLD – associated with a specific, on alcohol related problem such as drugs or toxins.

Toxin- associated steatohepatitis: associated with specific toxin or medication.eg petrochemical exposure in oil industry

CRITERIA FOR NAFLD

- 1) A liver histology showing moderate to gross macro vesicular fatty change with or without inflammation (lobular or portal), and Mallory bodies, fibrosis, or cirrhosis.*
- 1) No alcohol consumption or consumption less than 40 g of ethanol per week.**
- 2) Absence of serologic evidence of hepatitis B or hepatitis C.**

EPIDEMIOLOGY AND PREVALENCE

Prevalence of NAFLD 13-18% and that of NASH specifically 2-3% (1.2-9%) in two Japanese studies- incidence rate of 31 and 86 cases of suspected NAFLD person-years respectively. (AGA guidelines 2012)

The prevalence by USG is-46% and the prevalence by histology -12.2 %. (AGA guidelines 2012)

It is a disease of both the sexes, ethnicities, and all age groups (peak 40-59) higher in Hispanics than Africa Americans. Occurs more frequently in females (65 to 83%)

OBESITY

4.6 Fold increase risk of fatty liver in USG compared to non obese. Autopsy studies, steatosis-approx 70%obese vs. 35% lean. Severe obesity patients- prevalence of NAFLD may go up to 90%. Obese patients with abnormal liver enzymes approx 30% have sepal fibrosis and 10% cirrhosis.

INSULIN RESISTANCE AND T2 DM

Insulin resistance is very common in NAFLD. Progression to overt diabetes is preceded by steatosis in susceptible population. 75% of T2 DM- fatty infiltration. Liver injury worsens the preexistent diabetes in patients with NAFLD and doubles the prevalence of cirrhosis from 10-25%.

HYPERLIPIDEMIA

2/3 of hypertriglyceridemia&1/3rd with hypercholesterolemia- fatty liver Hyperlipidemia reported in 92% of NASH patients. Ethnic variation in lipid metabolism. Heterogenicity in lipid phenotypes in NAFLD

METABOLIC SYNDROME

Steatosis and central adiposity with underlying insulin resistance and lipotoxicity- independent risk factor for metabolic syndrome. Unexplained elevation of liver enzymes attributable to NAFLD seen in 7% of individuals with metabolic syndrome defined by adult Treatment Panel 111 criteria.

GENETIC AND FAMILIAL FACTORS

Ethnic variation: Hispanic descent in USA has higher prevalence of NASH than primarily african-american descent. It reflects the ethnic difference in body fat distribution and lipoprotein metabolism. Familial factors: High prevalence in 1st degree relatives in NASH. Insulin resistance among relatives of T2DM and impaired skeletal muscle mitochondrial metabolism in offspring supports genetic component. Genetic variation: gene coding phospholipase like protein called PNPLA3 or adiponutrin – major predictor of steatosis in different ethnic groups.

NASH in other liver diseases

Steatosis mediated by core protein metabolism and microsomal TGL transfer protein associated with hepatitis C {geno3} infection. Obesity, insulin resistance negatively influences response to antiviral therapy. Occult hemochromatosis and iron loading – cofactor in progression of NASH. Steatosis –potential factor in progression of PBC.

ETIOLOGY

Causes of Nonalcoholic Fatty Liver Disease

Acquired Metabolic Disorders

- Diabetes mellitus
- Dyslipidemia
- Kwashiorkor and marasmus
- Obesity
- Starvation

Cytotoxic and Cytostatic Drugs

Other Drugs and Toxins

- Amiodarone
- 4,4'-diethylaminoethoxyhexestrol
- Dichlorethylene
- Ethionine
- Ethyl bromide
- Estrogens
- Glucocorticoids
- Highly active antiretroviral therapy
- Hydrazine
- Hypoglycin

- Orotate
- Perhexilene maleate
- Safrole
- Tamoxifen

Acquired Metabolic Disorders

- Diabetes mellitus
- Dyslipidemia
- Kwashiorkor and marasmus
- Obesity
- Starvation

Acquired Metabolic Disorders

- Diabetes mellitus
- Dyslipidemia
- Kwashiorkor and marasmus
- Obesity
- Starvation

Cytotoxic and Cytostatic Drugs

Other Drugs and Toxins

- Amiodarone
- 4,4'-diethylaminoethoxyhexestrol
- Dichlorethylene

- Ethionine
- Ethyl bromide
- Estrogens
- Glucocorticoids
- Highly active antiretroviral therapy
- Hydrazine
- Hypoglycin
- Orotate
- Perhexilene maleate
- Safrole
- Tamoxifen

Metals

- Rare earths of low atomic number
- Thallium compounds
- Uranium compounds

Inborn Errors of Metabolism

- Abetalipoproteinemia
- Familial hepatosteatorrhea
- Galactosemia
- Glycogen storage disease
- Homocystinuria

- Systemic carnitine deficiency
- Tyrosinemia
- Weber-Christian syndrome
- Wilson disease


Surgical Procedures

- Biliopancreatic diversion
- Extensive small bowel resection
- Gastric and
- Jejunioileal bypass

Miscellaneous Conditions

- Industrial exposure to petrochemicals
- Inflammatory bowel disease
- Partial lipodystrophy
- Jejunal diverticulosis with bacterial overgrowth
- Severe anemia
- Total parenteral nutrition

PATHOGENESIS OF NAFLD



TRIGLYCERIDE ACCUMULATION

Occurs due to shifting of fatty acid metabolism to lipogenesis and also synthesis of lipoprotein decreases

- Excessive importation of FFA

Obesity

- Impaired VLDL synthesis and secretion
- Impaired beta-oxidation of FFA to ATP

INSULIN RESISTANCE& HYPERINSULINEMIA

It is a Primary pathogenic factor in steatosis. Excess FFA causes insulin resistance by down regulating IRS-1 signaling. Insulin resistance potentiated by aberrant function of peptide mediators TNF- α , leptin and adiponectin. TNF- α downregulates IRS-1 signaling via serine phosphorylation through activation of Jun terminal N kinase . Activation of inhibitor Kappa β kinase / nuclear factor kappa β by FFA – reduce insulin sensitivity.

LIPID PEROXIDATION & HEPATIC LIPOTOXICITY

FFA upregulates cyst P-450 – enhanced generation of ROS & lipid per oxidation. Increase FFA concentration –sustained upregulation of PPAR- α -promotes fatty acid oxidation. FFA – direct toxicity to cell membrane- toxic fatty acid ethylesters formation – disruption of mitochondrial function. Adiponectin and Leptin

Other mechanisms

- Portal endotoxemia
- Oxidative stress- induction of cyt P450 2E1- generate ROS – peroxidase cellular membrane- cell injury.
- Mitochondrial changes
 - Mitochondrial ROS formation
 - Megamitochondria and crystalline inclusions
 - Structural abnormalities
 - Altered ATP homeostasis

Clinical features


Clinical and Laboratory Features of Nonalcoholic Fatty Liver Disease

SYMPTOMS	SIGNS	LAB FEATURES
Common		
None (48-100%) Fatigue (70%)	Hepatomegaly	2 to 4 fold elevation of ALT/AST levels
RUQ pain (50%)		AST/ALT ratio <1
	Acantosis nigricans	S.ALP slightly elevated
	Occasional Neurological Deficits. In children	Normal bilirubin and albumin&PT levels Elevated serum ferritin levels. Increase in uric acid levels
Uncommon		
Vague right upper quadrant pain	Splenomegaly	Low titre<1:320)ANA positivity
Fatigue	Spider angiomas	Elevated transferrin saturation
Malaise	Palmar erythema	HFE gene mutation (C282Y)
	Ascites	Elevation serum Ig A

NAFLD and Normal Aminotranferases


- Occurs in later stages of steato hepatitis.
- “True normal” in obese patients may be lower than that of lean individuals.
- ALT levels are positively correlated with central obesity and hyperinsulinemia(leading to ongoing effort to revise normal changes).
- Glitazones normalise transminase levels,but stil criteria for NASH met on follow up liver biopsy.

Histopathology of NAFLD




- Picture of simple steatosis or fatty liver.
- Glycogenated nuclei seen.

NASH




- “Lobular inflammation” is hallmark.
- Hepatocyte ballooning and necrosis of varying degrees present.

NASH with cirrhosis



- NASH with cirrhosis (macrovesicular steatosis,inflammation and cirrhosis)
- Early Cirrhosis (bridging fibrosis)
- Masson trichrome staining.



- Mallory – Denker bodies – perinuclear material- dark staining brown masses.

Histological Features of NAFLD

Present in All or Most Cases

- Macrovesicular steatosis
- Diffuse or centrilobular steatosis; degree may correlate with BMI
- Parenchymal inflammation
- Polymorphonuclear neutrophils, lymphocytes, other mononuclear cells
- Hepatocyte necrosis
- Ballooning hepatocyte degeneration

Observed with Varied Frequencies

- Perivenular, perisinusoidal, or periportal fibrosis (37%-84%), moderate to severe in 15%-50%; most prevalent in zone 3 (perivenular)
- Cirrhosis (7%-16% on index biopsy specimen)
- Mallory bodies
- Glycogenated nuclei
- Lipogranulomas
- Stainable hepatic iron

Risk Factors for Advanced Nonalcoholic Fatty Liver Disease

Clinical

- Older age (>50 years)
- Obesity
- Diabetes mellitus/insulin resistance
- Hypertension

Laboratory

- AST/ALT ratio > 1
- Serum ALT level > twice the upper limit of normal
- Serum triglyceride levels > 155 mg/Dl

Histologic

- Severe steatosis
- Necroinflammatory activity (hepatocyte ballooning, necrosis)
- Stainable iron

Diagnostic approach and evaluation of NAFLD.

Essential Recording

- Age, Gender
- Body weight, body height, waist circumference, hip circumference, BMI
- Alcohol consumption (amount/frequency, at least g/day)

- History of liver disease, including HBV, HCV, autoimmune disease, DM, hyperlipidimia
- Family History of above mentioned (at least first degree)

NAFLD and Related Biochemical Studies Including OGTT and, IR


- Arthrometry: BMI, Wait/hip ratio
- AST/ALT, rGT, Bilirubin, ALP
- Cholesterol, HDL, TG, UA
- SBP/DBP
- OGTT
- Insulin Resistance
- ANA, Ferritin

HOMA & QUICKI

The screenshot shows a window titled "HOMA2 Calculator". Under the heading "Fasting values", there are two input fields: "Plasma glucose :" with a value of 7.8 and a unit selector set to mmol/l (with mg/dl as an alternative), and "Insulin" with a value of 65 and a unit selector set to pmol/l (with µU/ml as an alternative). Below these, the calculated results are displayed: %B : 45.6, %S : 74.5, and IR : 1.3. At the bottom, there are four buttons: "Calculate", "Copy", "Print", and "Exit".

- Used to estimate insulin resistance by utilising mathematical manipulation of fasting insulin and glucose levels.
- Homa index: $\text{insulin concentration } (\mu\text{U/ml}) \times \text{fasting glucose concentration (mmol/l)} / 22.5$ - utilise
- Cutoff level: 2.5 for adult
- Cutoff level: 3.16 for adolescent
 - Keskin M, et al. Padiatric 2005 March
 - QUICKI (quantitative insulin sensitivity index)- the addition of free fatty acid level in this will improve the accuracy

Diagnostic approach in suspected NAFLD



Role of Liver biopsy

The only diagnostic test that quantifies hepatic steatosis, necrosis and fibrosis and histological stage of NAFLD is best prognostic indicator.

- Indications in

NAFLD:

Peripheral stigmata of chronic liver disease

Splenomegaly

Cytopenia

Abnormal iron studies

Diabetes and/or significant obesity in an individual over the age of 45

BIOPSY SCORING

- NASH grade and stage (brunt) 0-8
- NASH activity index (NAI)- 0-12 accounts for steatosis, necroinflammatory and hepatocyte injury

NAFLD activity score (NAS)-0-8, steatosis 0-3, lobular and portal inflammation 0-3 cellular ballooning 0-2

NAFLD—Natural History

- Steatosis generally follows a benign course
- Steatosis can progress to NASH \pm fibrosis
- NASH with fibrosis has increased liver-related morbidity and mortality
- Olmsted large study 420pts with definite NAFLD on imaging and biopsy findings

cirrhosis developed in 3% of patients.

- 132 patients of NAFLD followed for 18yrs.

clinical outcomes based on degree of injury on an index liver biopsy specimen.

Cirrhosis and liver related death are common in NAFLD types 3 and 4 than 1 and 2.

- Studies show that long term survival of patients with NASH significantly better than alcoholic hepatitis .
- 5-10yr outcome of NAFLD associated cirrhosis was similar to that for HCV associated cirrhosis, although HCC less common in patients with NAFLD.

- Studies show that long term survival of patients with NASH significantly better than alcoholic hepatitis .

Independent predictors of fibrosis progression

- Diabetes mellitus,
- Low initial fibrosis stage
- Higher body mass index.
- Elevated liver enzymes

Non invasive markers of fibrosis in NAFLD

- Fibrotest: This tests incorporates haptoglobin,bil,GGTP,apo lipoproteinemiaA-1, α 2 macroglobulin and necroinflammatory index combines above markers plus ALT levels. cut off value – 0.70 has pos predictive value
- NAFLD fibrosis score : incorporates age ,BMI,hyperglycemia, AST/ALT ratio, platelet count and serum albumin level. PPV of 82-90%
- Transient elastography (fibro scan)
- S.dehydroepiandrosterone
- S.hyaluronic acid levels.
- Recently cytokeratin -18 level – novel biomarker for NASH

IMAGING OF NAFLD

- USG abdomen - “bright “ liver of increased echogenicity.

Poor detection if the degree of steatosis is less than 20% to 30%

Initial test of choice for large population screening.



- **CT imaging :**

Sensitivity and specificity of detecting fatty liver (with spleen-minus-liver attenuation of 10 Hounsfield units) were 0.84 and 0.99.
liver: spleen ratio <1- steatosis



- **MRI :**

T 1weighted image shows bright liver.


Dixon technique provides qualitative assessment of TGL content.

- **MR proton spectroscopy :**

Newer technique of quantification of fatty infiltration. most accurate method of quantifying steatosis



MR spectroscopy of TGL content in different ethnics



Currently non-invasive modalities are unable to detect NASH with or without fibrosis

MATERIALS AND METHODS

MATERIAL AND METHODS

The study was conducted at the department of Medical Gastroenterology, Institute of Pathology and in the Bernard Institute of Radiology Department in the Rajiv Gandhi Government General Hospital Chennai. and in the Madras Medical College Chennai.,

Patients coming to our OPD undergo routine investigations like CBC,LFT,RFT,Blood sugar and USG Abdomen, among them those who had fatty liver on USG abdomen were selected, those who with history of alcohol use and viral hepatitis were excluded

Inclusion criteria	1. Fatty liver on USG Abdomen
Exclusion criteria	<ol style="list-style-type: none">1. Alcohol (more than 20 g/d)2. HCV infection3. HBV infection4. Decompensated CLD5. HCC / focal lesion on USG or CT6. Coagulopathy (for liver Bx)7. Secondary cause- surgery, Drugs, Pregnancy

Total no of patients selected from the study is 30

Table 1
SEX DISTRIBUTION

SEX DISTRIBUTION	CASES
MALE	15
FEMALE	15

SEX DISTRIBUTION




Table 2

AGE DISTRIBUTION

AGE DISTRIBUTION	MALE	FEMALE	TOTAL
20-30	1	0	1
30-40	4	3	7
40-50	3	10	13
50-60	7	2	9

AGE DISTRIBUTION





Table 3
BMI CHART

BMI	MALE	FEMALE	TOTAL
< 20	4	3	7
20-25	3	3	6
25-30	6	8	14
30-35	2	1	3

BMI CHART



These selected patient subjected to the following additional investigations Biochemical investigation – LFT. HBsAg, anti HCV, BT, CT, PT/INR.

Detailed history was taken and physical examinations done. All the patients underwent non contrast CT [ncCT] scan. those patients underwent non contrast computerized tomography of the abdomen with machine [GE Medical Systems, Milwaukee, WI] of Bernard Institute Of Radiology in Rajiv Gandhi Government General Hospital – Chennai .

Following observation are assessed from the Radiology Department. With Non contrast computerized tomography [nc CT] CT attenuation index in houns field units [HU] of liver and CT attenuation index in houns field units [HU] of spleen were assessed.

All the patients after informed concern underwent liver biopsy after proper aseptic precaution and after screening for coagulation profile in the side room of ward no : 243 and 245 [male and female ward] of Medical Gastro Enterology Department Liver biopsy was done with biopsy true cut biopsy needle using biopsy gun, the specimen was placed in 10% formalin and submitted for histologic examination.

LIVER BIOPSY GUN



After the liver biopsy all the patients were admitted and observed in the ward for 24 hrs. Successful biopsy specimens were obtained in all the patients among them two patients [mrs.Ayisha bevi and mr.Prasanth] required two pass for adequate biopsy sample size, other 28 patients required one pass only.

Among them one patient [mrs.uniammal] developed severe pain after the procedure [pain radiating to right shoulder and became dyspneic] she has been screened with X ray chest, X ray erect abdomen and USG Abdomen .no evidence of any pneumothorax noted and the patient was managed symptomatically . no intervention required and sent home after 48 hrs of observation.

RESULTS

RESULTS

This study includes total of 30 cases- CT Finding of six patients




Mrs.Muniammal



Mrs.Nisha



Mr.Prasanth



Mrs. Priya



Mr.Parthiban




Mr.Vetri selvan

Table 4**CT Scan Finding of 30 patients –Attenuation in HOUNS FIELD****UNITS [HU]**

NAME	AGE	SEX	CT HU Liver	CT HU spleen	CT HU Liver- spleen
Raja	29	M	26	49	-23
Banumathi	32	F	31	52	-21
Devaki	60	F	26	45	-19
Rangasamy	55	M	28	47	-19
Yoganand	57	M	45	54	-9
Aisha bee	47	F	34	42	-8
Purushothaman	39	M	36	48	-12
Ladly	32	M	26	48	-22
Ram mohan	53	M	26	49	-23
Prabha	44	F	26	48	-22
Ramasamy	55	M	26	42	-6
Ayisha bevi	47	F	28	45	-17
Prasanth	44	M	29	54	-25
Jananki	39	F	26	42	-18
Gopi	43	M	28	48	-20
Nisha	47	F	26	57	-25
Priya	34	F	34	51	-17
Jothi	42	F	28	45	-17
Amudha	48	F	28	45	-16
Lalitha	51	F	28	45	-17
Kumar	45	M	31	45	-14
Muniammal	48	M	26	45	-19

Yogas	57	M	32	45	-13
Vetriselvan	35	M	26	56	-24
Vijayalakshmi	39	F	29	45	-16
Sangeetha	40	F	29	48	-19
Noorjagans	42	F	26	48	-19
Anandvalli	45	F	28	45	-17
Thiru	51	M	26	48	-22
Parthiban	54	M	28	50	-22



Among them their biopsy has been graded and its results


Table 5 (a)
BIOPSY GRADING ADOPTED

Grade 0	Minimal <5 % steatosis
Grade I	5-33% steatosis
Grade II	33-66 % steatosis
Grade III	>66 % steatosis



Table 5 (b)
HISTOLOGICAL GRADE-OBSERVATIONS

HISTOLOGIC GRADE	MALE	FEMALE	TOTAL
MINIMAL	2	3	5
GRADE I	5	5	10
GRADE II	4	7	11
GRADE III	3	1	4

HISTOLOGICAL GRADE-OBSERVATIONS-DIAGRAM





Minimal Macro-Steatosis





Grade I

Macro-Steatosis



Grade II

Macro-Steatosis



Grade III

Macro-Steatosis




Table 6**One way ANOVA to compare mean values between Histology**

	Histology	N	Mean	Std. Dev	Min	Max	P-Value
CT HU Liver-spleen	Minimal	5	13.00	4.36	8.00	18.00	0.001
	Mild	9	15.11	4.17	6.00	19.00	
	Moderate	12	20.25	2.05	17.00	23.00	
	Severe	4	24.25	0.96	23.00	25.00	
	Total	30	18.03	4.85	6.00	25.00	

Table 7**Tukey HSD Post Hoc Tests to compare multiple pairwise comparisons**

Dependent Variable	Pairs		Mean Difference	P-Value
CT HU Liver-spleen	Minimal	Mild	-2.111	0.640
		Moderate	-7.250	0.001
		Severe	-11.250	0.001
	Mild	Moderate	-5.139	0.006
		Severe	-9.139	0.001
	Moderate	Severe	-4.000	0.157





Table 8

One way ANOVA to compare mean values between Histology

	Histology	N	Mean	Std. Dev	Min	Max	P-Value
Liver Spleen Ratio	Minimal	5	0.719	0.097	0.620	0.830	0.001
	Mild	9	0.638	0.056	0.580	0.750	
	Moderate	12	0.576	0.041	0.530	0.670	
	Severe	4	0.497	0.043	0.460	0.540	
	Total	30	0.608	0.087	0.460	0.830	



Tukey HSD Post Hoc Tests to compare multiple pairwise comparisons

Dependent Variable	Pairs		Mean Difference	P-Value
Liver Spleen Ratio	Minimal	Mild	0.081	0.081
		Moderate	0.143	0.001
		Severe	0.222	0.001
	Mild	Moderate	0.062	0.092
		Severe	0.141	0.002
	Moderate	Severe	0.079	0.110

Tukey HSD Post Hoc Tests to compare multiple pairwise comparisons

Dependent Variable	Pairs		Mean Difference	P-Value
CT HU Liver	Minimal	Mild	4.111	0.233
		Moderate	5.167	0.072
		Severe	6.250	0.090
	Mild	Moderate	1.056	0.920
		Severe	2.139	0.783
	Moderate	Severe	1.083	0.959

Tukey HSD Post Hoc Tests to compare multiple pairwise comparisons


Dependent Variable	Pairs		Mean Difference	P-Value
BMI	Minimal	Mild	-0.249	1.000
		Moderate	1.173	0.967
		Severe	-1.660	0.954
	Mild	Moderate	1.422	0.905
		Severe	-1.411	0.960
	Moderate	Severe	-2.833	0.734

Tukey HSD Post Hoc Tests to compare multiple pairwise comparisons

Variable	Pairs		Mean Difference	P-Value
Liver Spleen Ratio	Mild	Moderate	.05064	0.280
		Severe	.06243	0.608
	Moderate	Severe	.01180	0.982

CHARTS

Correlation Graph



DISCUSSION

DISCUSSION

The degree of steatosis in NAFLD patients is usually assessed with invasive technique like liver biopsy [Leon A. Adams, et al]⁷. Because of the invasive method of the liver biopsy and its painful nature of the procedure with moderate complication rate there has always been a need to find alternative less invasive, less painful, quick method to accurately diagnose NAFLD is the need of the hour.

CT scan and MRI scan are well established imaging modalities for diagnosis of various disorders. These imaging modalities can also assess the degree of fat as accurately as histology of liver tissue. This accuracy of grading of fatty liver by CT and MRI has been utilized by various investigators. Olmsted et al large study 420pts with definite NAFLD on imaging and liver biopsy. But only few studies are available [Nathan et al; sheela ramesh; arun j.sanyal et al]^{1,2,3}.

These authors tried to validate CT scan as non invasive diagnostic modality for this purpose, by comparing against gold standard liver biopsy. but their studies have been done in small number [ie., 15 subjects].

Most of the studies are with magnetic resonance image [MRI]. few studies are with computerized tomography [Nathan et al]¹. These studies are also small in number.

In our study non contrast computerized tomography scan [nc CT] has been done in all the patients [no of sample size is 30] compared to previous studies the number is doubled.

Initial evaluation of NAFLD requires ultra sound scan of abdomen. just as In previous studies the investigators used USG for initial evaluation. But this USG abdomen scan can be used for screening purpose only. It is not useful for semi quantitative assessment of steatosis.

Non contrast CT scan [nc CT] has been done in both the studies. Because non contrast CT scan will give accurate measurement of fat content of the liver measured as a hounce field unit is compared with nearby organ spleen. Both the studies [our study and the previous study] followed the same method. Use of contrast is avoided because contrast will decrease the liver attenuation in hounce field unit. It will give false positive results.

The diagnostic sensitivity of the CT depends upon the severity of steatosis. HU will fall when more steatosis is present in the liver. In the milder form of steatosis the HU will be in higher values or equal to spleen value.

The study has confirmed the nc CT is useful technique delineating macrosteatosis (HMS) in very high BMI P&S , CT with more HU unit

difference between CTHU liver to spleen is significantly and directly correlated to the histology in HMS.

CT assessment of hepatic steatosis in previous studies was criticized on the basis of limited accuracy to estimate hepatic fat content accurately [etal] but such criticism cannot be leveled with our data

Most of the studies conducted earlier [Nathan j.shore et al]¹ is not a direct comparison between macrosteatosis to liver HU unit difference, only one study compared [Nathan et al]¹ hepatic macrosteatosis HMS to CT HU – liver difference, those studies are in small number.

The strength of this study, that it is a direct comparison of liver CT attenuation with histological grade of steatosis, it is not included the stage of fibrosis. In our study we have adopted grading of steatosis in four stages namely grade 0 is minimal <5 % of macrosteatosis, grade I is 5-33 % of macrosteatosis, grade II is 33-66 % of macrosteatosis, and grade III is more than 66 % When compared to previous studies [Nathan j.shore et al]¹ adapted hepatic macrosteatosis as more than 30 % [HMS]. and this value has been compared to hepatic attenuation index in HU and also there is no control arm in the previous mentioned studies [Nathan j.shore et al]¹. To delineate in which grade of HMS will directly correlate to hepatic CT attenuation index.

Nathan j.shore etal¹ in their study have taken biopsy from both lobes of liver. And the biopsy is a wedge biopsy taken at the time of bariatric surgery. The tissue sample size is compared to our study is larger. And also they studied the liver tissue triglyceride's concentration by biochemical methods. But in our study biopsy has been taken by true cut needle biopsy using biopsy gun and the tissue sample is small and only taken from the right lobe of the liver alone, in this study no biochemical analysis of liver tissue is made ,the possibility of false negative results [if the normal liver tissue in the biopsied sample]could be possible and a false positive results [a focal nodular fatty liver] may be another possibility to give grade III macro steatosis on histology.

Previous study by qayyum etal. In his study he has included higher BMI patients only [BMI > 30] when compared to this study where in we have included all the patients with BMI starting from 17.5 - 32 . In our study the inclusion criteria is based on the ultra sonogram finding of fatty liver. So that lean NASH can be calculated with our study. Out of 30 patients lean NASH is [0 %].

In various studies sample size is small less than 15 patients [Nathan j.shore etal]¹compared to this study the sample size is 30 subjects. But one drawback of our study, there is no control arm, if a control arm is included we can accurately delineate at which liver HU difference which

we can accurately predict the presence of NAFLD. It could served as a quantitative assessment of NAFLD score. The control arm was not designed due to the ethical point of view as well as risk of liver biopsy in normal subjects. Even though if the normal subject found to be detected having NAFLD. There is no definite treatment option available till Date. Also associated risk of radiation exposure in normal subjects for an investigation for a non clinical purpose. The studied patients under went plain CT abdomen, there is a life time risk of medical radiation but is a modest radiation exposure and also the abdominal region only exposed to radiation and not to the entire body.

CONCLUSION

CONCLUSION

- 1] There is definite correlation between non contrast computerized tomography [nc CT] with liver biopsy in this study [p value 0.001]
- 2] Hence non contrast computerized tomography [nc CT] can be used as a sole investigation for defining NAFLD. invasive liver biopsy can be used for borderline or indeterminate cases

BIBLIOGRAPHY

REFERENCES

1. Nathan J sores, Flegal KM, Carroll MD, Ogden CL, Johnson CL. Prevalence and trends in obesity among US adults, 1999–2000. JAMA. 2002;288:1723–1727. doi: 10.1001/jama.288.14.1723. [\[PubMed\]](#) [\[Cross Ref\]](#)
2. Nair S, Mason A, Eason J, Loss G, Perrillo RP. Is obesity an independent risk factor for hepatocellular carcinoma in cirrhosis? Hepatology. 2002;36:150–155. doi: 10.1053/jhep.2002.33713. [\[PubMed\]](#) [\[Cross Ref\]](#)
3. Wanless IR, Lentz JS. Fatty liver hepatitis (steatohepatitis) and obesity: An autopsy study with analysis of risk factors. Hepatology. 1990;12:1106–1110. doi: 10.1002/hep.1840120505. [\[PubMed\]](#) [\[Cross Ref\]](#)
4. Adinolfi LE, Gambardella M, Andreana A, Tripodi MF, Utili R, Ruggiero G. Steatosis accelerates the progression of liver damage of chronic hepatitis C patients and correlates with specific HCV genotype and visceral obesity. Hepatology. 2001;33:1358–1364. doi: 10.1053/jhep.2001.24432. [\[PubMed\]](#) [\[Cross Ref\]](#)
5. Browning JD, Szczepaniak LS, Dobbins R, Nuremberg P, Horton JD, Cohen JC, Grundy SM, Hobbs HH. Prevalence of hepatic steatosis in an urban population in the United States: Impact of ethnicity. Hepatology. 2004;40:1387–1395. doi: 10.1002/hep.20466. [\[PubMed\]](#) [\[Cross Ref\]](#)
6. Marchesini G, Bugianesi E, Forlani G, Cerrelli F, Lenzi M, Manini R, Natale S, Vanni E, Villanova N, Melchionda N, Rizzetto M.

Nonalcoholic fatty liver, steatohepatitis, and the metabolic syndrome. *Hepatology*. 2003;37:917â€“923. doi: 10.1053/jhep.2003.50161. [[PubMed](#)] [[Cross Ref](#)]

7. Xirouchakis E, Sigalas A, Manousou P, Calvaruso V, Pleguezuelo M, Corbani A, Maimone S, Patch D, Burroughs AK. Models for non-alcoholic fatty liver disease: A link with vascular risk. *Curr Pharm Des*. 2008;14:378â€“384. doi: 10.2174/138161208783497705. [[PubMed](#)] [[Cross Ref](#)]
8. Rockey DC, Caldwell SH, Goodman ZD, Nelson RC, Smith AD. Liver biopsy. *Hepatology*. 2009;49:1017â€“1044. doi: 10.1002/hep.22742. [[PubMed](#)] [[Cross Ref](#)]
9. Qayyum A, Chen DM, Breiman RS, Westphalen AC, Yeh BM, Jones KD, Lu Y, Coakley FV, Callen PW. Evaluation of diffuse liver steatosis by ultrasound, computed tomography, and magnetic resonance imaging: Which modality is best? *Clin Imaging*. 2009;33:110â€“115. doi: 10.1016/j.clinimag.2008.06.036. [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]
10. Schwenzer NF, Springer F, Schraml C, Stefan N, Machann J, Schick F. Non-invasive assessment and quantification of liver steatosis by ultrasound, computed tomography and magnetic resonance. *J Hepatol*. 2009;51:433â€“445. doi: 10.1016/j.jhep.2009.05.023. [[PubMed](#)] [[Cross Ref](#)]
11. Szczepaniak LS, Babcock EE, Schick F, Dobbins RL, Garg A, Burns DK, McGarry JD, Stein DT. Measurement of intracellular triglyceride stores by H spectroscopy: Validation in vivo. *Am J Physiol*. 1999;276:E977â€“E989. [[PubMed](#)]

12. Vuppalanchi R, Cummings OW, Saxena R, Ulbright TM, Martis N, Jones DR, Bansal N, Chalasani N. Relationship among histologic, radiologic, and biochemical assessments of hepatic steatosis: A study of human liver samples. *J Clin Gastroenterol*. 2007;41:206–210. doi: 10.1097/01.mcg.0000225515.28536.3a. [\[PubMed\]](#) [\[Cross Ref\]](#)
13. Lee SW, Park SH, Kim KW, Choi EK, Shin YM, Kim PN, Lee KH, Yu ES, Hwang S, Lee SG. Unenhanced CT for assessment of macrovesicular hepatic steatosis in living liver donors: Comparison of visual grading with liver attenuation index. *Radiology*. 2007;244:479–485. doi: 10.1148/radiol.2442061177. [\[PubMed\]](#) [\[Cross Ref\]](#)
14. Contos MJ, Sanyal AJ. The clinicopathologic spectrum and management of nonalcoholic fatty liver disease. *Adv Anat Pathol* 2002;9: 37–51.
15. Clark JM, Brancati FL, Diehl AM. The prevalence and etiology of elevated aminotransferase levels in the United States. *Am J Gastroenterol* 2003;98:960–967.
16. Third Report of the National Cholesterol Education Program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (adult treatment panel III) final report. *Circulation* 2002;106:3143–3421.
17. Sanyal AJ, Campbell-Sargent C, Mirshahi F, Rizzo WB, Contos MJ, Sterling RK, et al. Nonalcoholic steatohepatitis: association of insulin resistance and mitochondrial abnormalities. *Gastroenterology* 2001; 120:1183–1192.

18. Chitturi S, Farrell GC. Etiopathogenesis of nonalcoholic steatohepatitis. *Semin Liver Dis* 2001;21:27–41.
 19. Ford ES, Giles WH, Dietz WH. Prevalence of the metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. *JAMA* 2002;287:356–359.
 20. Flegal KM, Carroll MD, Ogden CL, Johnson CL. Prevalence and trends in obesity among US adults, 1999–2000. *JAMA* 2002;288:1723–1727.
 21. Haque M, Sanyal AJ. The metabolic abnormalities associated with non-alcoholic fatty liver disease. *Best Pract Res Clin Gastroenterol* 2002;16:709–731.
 22. Skelly MM, James PD, Ryder SD. Findings on liver biopsy to investigate abnormal liver function tests in the absence of diagnostic serology. *J Hepatol* 2001;35:195–199.
 23. Mathiesen UL, Franzen LE, Fryden A, Foberg U, Bodemar G. The clinical significance of slightly to moderately increased liver transaminase values in asymptomatic patients. *Scand J Gastroenterol* 1999;34:85–91.
 24. Choudhury J, Sanyal AJ. Insulin resistance and the pathogenesis of nonalcoholic fatty liver disease. *Clin Liver Dis* 2004;8:575–594.
- [12] De Fronzo RA, Ferrannini E. Insulin resistance. A multifaceted syndrome responsible for NIDDM, obesity, hypertension, dyslipidemia, and atherosclerotic cardiovascular disease. *Diab Care* 1991;14: 173–194.

25. Diehl AM. Fatty liver, hypertension, and the metabolic syndrome. *Gut* 2004;53:923–924.
26. Marchesini G, Bugianesi E, Forlani G, Cerrelli F, Lenzi M, Manini R, et al. Nonalcoholic fatty liver, steatohepatitis, and the metabolic syndrome. *Hepatology* 2003;37:917–923.
27. Clark JM, Diehl AM. Hepatic steatosis and type 2 diabetes mellitus. *Curr Diab Rep* 2002;2:210–215.
28. Falck-Ytter Y, Younossi ZM, Marchesini G, McCullough AJ. Clinical features and natural history of nonalcoholic steatosis syndromes. *Semin Liver Dis* 2001;21:17–26.
29. Youssef W, McCullough AJ. Diabetes mellitus, obesity, and hepatic steatosis. *Semin Gastrointest Dis* 2002;13:17–30.
30. Neuschwander-Tetri BA, Bacon BR. Nonalcoholic steatohepatitis. *Med Clin North Am* 1996;80:1147–1166.
31. Haller DL, Sargeant C, Luketic VA, Sanyal AJ. A behavioral analysis of nonalcoholic fatty liver disease. *Hepatology* 2003;38:233A.
32. Bacon BR, Farahvash MJ, Janney CG, Neuschwander-Tetri BA. Nonalcoholic steatohepatitis: an expanded clinical entity. *Gastroenterology* 1994;107:1103–1109.
33. Ludwig J, Viggiano TR, McGill DB, Oh BJ. Nonalcoholic steatohepatitis: Mayo Clinic experiences with a hitherto unnamed disease. *Mayo Clin Proc* 1980;55:434–438.

34. Powell EE, Cooksley WG, Hanson R, Searle J, Halliday JW, Powell LW. The natural history of nonalcoholic steatohepatitis: a follow-up study of forty-two patients for up to 21 years. *Hepatology* 1990;11:74–80.
35. S. Ramesh, A.J. Sanyal / *Journal of Hepatology* 42 (2005) S2–S12 S11 [23] Clouston AD, Powell EE. Interaction of non-alcoholic fatty liver disease with other liver diseases. *Best Pract Res Clin Gastroenterol* 2002;16:767–781.
36. Matteoni CA, Younossi ZM, Gramlich T, Boparai N, Liu YC, McCullough AJ. Nonalcoholic fatty liver disease: a spectrum of clinical and pathological severity. *Gastroenterology* 1999;116:1413–1419.
37. Angulo P, Keach JC, Batts KP, Lindor KD. Independent predictors of liver fibrosis in patients with nonalcoholic steatohepatitis. *Hepatology* 1999;30:1356–1362.
38. Sanyal AJ. AGA technical review on nonalcoholic fatty liver disease. *Gastroenterology* 2002;123:1705–1725.
39. Yajima Y, Ohta K, Narui T, Abe R, Suzuki H, Ohtsuki M. Ultrasonographical diagnosis of fatty liver: significance of the liverkidney contrast. *Tohoku J Exp Med* 1983;139:43–50.
40. Fusamoto H, Suzuki K, Hayashi N, Sasaki Y, Kono M, Kasahara A, et al. Obesity and liver disease: evaluation of fatty infiltration of the liver using ultrasonic attenuation. *J Nutr Sci Vitaminol (Tokyo)* 1991; 37:S71–S77.

41. Lee RG. Nonalcoholic steatohepatitis: a study of 49 patients. *Hum Pathol* 1989;20:594–598.
42. Saadeh S, Younossi ZM, Remer EM, Gramlich T, Ong JP, Hurley M, et al. The utility of radiological imaging in nonalcoholic fatty liver disease. *Gastroenterology* 2002;123:745–750.
43. Jacobs JE, Birnbaum BA, Shapiro MA, Langlotz CP, Slosman F, Rubesin SE, et al. Diagnostic criteria for fatty infiltration of the liver on contrast-enhanced helical CT. *AJR Am J Roentgenol* 1998;171:659–664.
45. Sorbi D, McGill DB, Thistle JL, Therneau TM, Henry J, Lindor KD. An assessment of the role of liver biopsies in asymptomatic patients with chronic liver test abnormalities. *Am J Gastroenterol* 2000;95:3206–3210.
46. Caldwell SH, Oelsner DH, Iezzoni JC, Hespenheide EE, Battle EH, Driscoll CJ. Cryptogenic cirrhosis: clinical characterization and risk factors for underlying disease. *Hepatology* 1999;29:664–669.
47. Sorbi D, Boynton J, Lindor KD. The ratio of aspartate aminotransferase to alanine aminotransferase: potential value in differentiating nonalcoholic steatohepatitis from alcoholic liver disease. *Am J Gastroenterol* 1999;94:1018–1022.
48. Lewis KO, Paton A. ABC of alcohol: tools of detection. *Br Med J (Clin Res Ed)* 1981;283:1531–1532.
49. Salaspuro M. Use of enzymes for the diagnosis of alcohol-related organ damage. *Enzyme* 1987;37:87–107.

50. Nalpas B, Vassault A, Charpin S, Lacour B, Berthelot P. Serum mitochondrial aspartate aminotransferase as a marker of chronic alcoholism: diagnostic value and interpretation in a liver unit. *Hepatology* 1986;6:608–614.
51. Storey EL, Mack U, Powell LW, Halliday JW. Use of chromatofocusing to detect a transferrin variant in serum of alcoholic subjects. *Clin Chem* 1985;31:1543–1545.
52. Marchesini G, Brizi M, Bianchi G, Tomassetti S, Bugianesi E, Lenzi M, et al. Nonalcoholic fatty liver disease: a feature of the metabolic syndrome. *Diabetes* 2001;50:1844–1850.
53. Garg A, Misra A. Hepatic steatosis, insulin resistance, and adipose tissue disorders. *J Clin Endocrinol Metab* 2002;87:3019–3022.
54. Garg A. Lipodystrophies. *Am J Med* 2000;108:143–152.
55. Executive summary of the clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults. *Arch Intern Med* 1998;158:1855–1867.
56. Bray GA, Nielsen SJ, Popkin BM. Consumption of high-fructose corn syrup in beverages may play a role in the epidemic of obesity. *Am J Clin Nutr* 2004;79:537–543.
57. Archer WR, Lamarche B, Deriaz O, Landry N, Corneau L, Despres JP, et al. Variations in body composition and plasma lipids in response to a high-carbohydrate diet. *Obes Res* 2003;11:978–986.
58. Sondike SB, Copperman N, Jacobson MS. Effects of a lowcarbohydrate diet on weight loss and cardiovascular risk factor in overweight adolescents. *J Pediatr* 2003;142:253–258.

59. Hayes KC. Saturated fats and blood lipids: new slant on an old story. *Can J Cardiol* 1995;11:39G–446.
60. Kwiterovich Jr PO. The effect of dietary fat, antioxidants, and prooxidants on blood lipids, lipoproteins, and atherosclerosis. *J Am Diet Assoc* 1997;97:S31–S41.
61. Gaziano JM, Manson JE. Diet and heart disease. The role of fat, alcohol, and antioxidants. *Cardiol Clin* 1996;14:69–83.
62. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, et al. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393–403.
63. Kriska A. Can a physically active lifestyle prevent type 2 diabetes? *Exerc Sport Sci Rev* 2003;31:132–137.
64. Rollins G. Moderate exercise reduces the risk of heart disease and death in men with type 2 diabetes. *Rep Med Guidelines Outcomes Res* 2003;14:10–12.
65. Hollander P. Orlistat in the treatment of obesity. *Prim Care* 2003;30:427–440.
66. Holt R. Orlistat reduces features of the metabolic syndrome: the XENDOS study. *Diabetes Obes Metab* 2003;5:356.
67. Padwal R, Li SK, Lau DC. Long-term pharmacotherapy for overweight and obesity: a systematic review and meta-analysis of randomized controlled trials. *Int J Obes Relat Metab Disord* 2003;27: 1437–1446.

68. Ryan DH. Use of sibutramine to treat obesity. *Prim Care* 2003;30: 405–426.
69. Sabuncu T, Nazligul Y, Karaoglanoglu M, Ucar E, Kilic FB. The effects of sibutramine and orlistat on the ultrasonographic findings, insulin resistance and liver enzyme levels in obese patients with nonalcoholic steatohepatitis. *Rom J Gastroenterol* 2003;12:189–192.
70. Luyckx FH, Scheen AJ, Desaive C, Thiry A, Lefebvre PJ. Parallel reversibility of biological markers of the metabolic syndrome and liver steatosis after gastroplasty-induced weight loss in severe obesity. *J Clin Endocrinol Metab* 1999;84:4293.
71. Luyckx FH, Desaive C, Thiry A, Dewe W, Scheen AJ, Gielen JE, et al. Liver abnormalities in severely obese subjects: effect of drastic weight loss after gastroplasty. *Int J Obes Relat Metab Disord* 1998;22: 222–226.
72. Caldwell SH, Hespenheide EE, Redick JA, Iezzoni JC, Battle EH, Sheppard BL, et al. A pilot study of a thiazolidinedione, troglitazone, in nonalcoholic steatohepatitis. *Am J Gastroenterol* 2001;96:519–525.

ANNEXURES

CONSENT FORM

DESCRIPTION: You are invited to participate in a research study on liver disease. We are going to assess the normal values of liver biochemistry ,liver scan and liver tissue biopsy in healthy adults.

PROCEDURES: You will be asked to do ultra sound scan , CT-Scan, Liver biopsy (small piece of liver tissue by a small needle biopsy)and Blood test provide a sample of blood (1 tablespoon). The blood will be taken from your arm at the same time you are going to donate the blood.

The results of the study of your samples will be used for research purposes only and you will not be told the results of the tests.

RISKS AND BENEFITS: there are mild risks of bleeding from liver puncture site associated pain with this study.

TIME INVOLVEMENT: Post liver biopsy you have to stay for one day for observation of any complication

PAYMENTS: You will not be paid to participate in this study.

I, _____ , hereby consent to do USG abdomen, CT abdomen , liver biopsy ,Blood sample which I have donated as a voluntarily for the purpose of estimating the function of the liver in normal individuals. I have understood that this sample will be used for research and the results will not be available to me. I have also understood that this does not involve any additional procedure.

Witness – Signature, Name and address

Signature of the donor

Name of the Donor

PROFORMA

NAME	
AGE/SEX	
GE.No.	
CBC	
LFT	
PT/INR	
BT/CT	
HBsAg	
AntiHCV	
BLOOD SUGAR F/PP	
BMI WAIST/HIP	
ALOHOL CONSUPTION QUANTITY IN GRAMS	
USG ABDOMEN	
CT ABDOMEN LIVER/SPLEEN RATIO LIVER-SPLEEN INDEX	
LIEVER BIOPSY LENGTH FINDING COPLICATION	

MASTER CHART


SL.No	NAME	AGE	SEX	HEIGHT	WEIGHT	BMI	USG abdomen	CT HU Liver	CT HU spleen	CT HU Liver-spleen	HISTOLOGY	LFT1	LFT-AST	LFT-ALT	LFT-SAP
1	raja	29	m	155	75	31.2	3	26	49	-23	3	1.12	25.7	39.9	131
2	banumathi	32	f	155	70	29.1	1	31	52	-21	2	1.4	29	32	120
3	devaki	60	f	152	68	29.4	2	26	45	-19	2	0.9	34	42	92
4	rangasamy	55	m	168	72	25.5	1	28	47	-19	1	0.9	30	23	83
5	yoganand	57	m	168	80	28.3	2	45	54	-9	0	1.1	23	29	92
6	aisha bee	47	f	150	72	32	1	34	42	-8	0	0.7	59	44	152
7	purushothaman	39	m	169	80	28	1	36	48	-12	1	0.9	36	39	75
8	ladly	32	m	152	70	30.3	2	26	48	-22	2	1.1	94	86	132
9	ram mohan	53	m	155	69	28.7	1	26	49	-23	2	1.3	21	26	102
10	prabha	44	f	153	45	19.2	2	26	48	-22	2	1	56	76	134
11	ramasamy	55	m	165	50	18.4	3	26	42	-6	1	0.7	34	36	120
12	ayisha bevi	47	f	156	69	28.4	1	28	45	-17	1	1.2	29	23	84
13	prasanth	44	m	160	65	25.4	2	29	54	-25	3	0.9	102	109	196
14	jananki	39	f	158	50	20	1	26	42	-18	0	0.7	34	31	98
15	gopi	43	m	162	52	19.8	1	28	48	-20	2	0.8	34	36	102
16	nisha	47	f	158	71	28.4	2	26	57	-25	3	4.1	32	34	98
17	priya	34	f	157	68	27.6	2	34	51	-17	2	0.7	231	246	163
18	jothi	42	f	154	48	20.2	2	28	45	-17	2	0.8	24	19	98
19	amudha	48	f	160	48	18.7	2	28	45	-16	1	2.1	19	17	102
20	lalitha	51	f	149	63	28.4	1	28	45	-17	1	0.8	83	96	159
21	kumar	45	m	162	54	20.6	2	31	45	-14	0	1.8	19	17	109
22	muniammal	48	m	158	70	28	1	26	45	-19	1	0.8	64	96	128
23	yogas	57	m	168	70	24.8	1	32	45	-13	1	0.9	124	34	102
24	vetriselvan	35	m	159	53	21	2	26	56	-24	3	1.8	34	31	103
25	vijayalakshmi	39	f	155	56	23.3	1	29	45	-16	0	0.7	96	96	104
26	sangeetha	40	f	154	62	26.1	2	29	48	-19	2	2.1	19	20	96
27	noorjagan	42	f	150	42	17.5	2	26	48	-19	2	0.9	47	51	102
28	anandvalli	45	f	143	68	25.6	1	28	45	-17	1	1.2	29	34	106
29	thiru	51	m	167	47	17.7	2	26	48	-22	2	1	34	39	123
30	parthiban	54	m	165	50	18.4	2	28	50	-22	2	1	29	34	96

Gmail: Email from Google

Turnitin

https://turnitin.com/s_class_portfolio.asp?r=13.0858314612443&svr=6&lang=en_us&aid=80345&cid=5807035

Senthilkumar P 16102505 D.M. Medical Gastroenterology | User Info | Messages | Student | English | What's New | Help | Logout



Class Portfolio

Peer Review

My Grades

Discussion




Calendar

NOW VIEWING: HOME > TNMGRMU APRIL 2013 EXAMINATIONS

Class Homepage

This is your class homepage. To submit to an assignment click on the "Submit" button to the right of the assignment name. If the Submit button is grayed out, no submissions can be made to the assignment. If resubmissions are allowed the submit button will read "Resubmit" after you make your first submission to the assignment. To view the paper you have submitted, click the "View" button. Once the assignment's post date has passed, you will also be able to view the feedback left on your paper by clicking the "View" button.

Assignment Inbox: TNMGRMU APRIL 2013 EXAMINATIONS

	Info	Dates	Similarity	
Medical		Start 21-Nov-2012 11:24AM Due 31-Mar-2013 11:59PM Post 01-Apr-2013 12:00AM	20% 	<div>ResubmitViewDownload</div>
Dental		Start 27-Nov-2012 12:43PM Due 31-Dec-2012 11:59PM Post 07-Jan-2013 12:00AM		<div>SubmitViewDownload</div>

start

Downloads

IMG_0006 - Windo...

Transcend (E:)

KINGSTON (F:)

Turnitin - Google C...

Task Manager

turnitin [Compatibili...

3:25 PM

Turnitin Document Viewer - Google Chrome

https://turnitin.com/dv?o=312342535&u=1017270583&s=8&student_user=18&lang=en_us

TNMGRMU APRIL 2013 EXAMINA...Medical - DUE 31-Mar-2013What's New

OriginalityGradeMarkPeerMark

"Non-contrasted Computed Tomography for the Accurate Measurement of
BY: SENTHILKUMAR P. 16102605 D.M. MEDICAL GASTROENTEROLOGY

turnitin

20%
SIMILAR

--
OUT OF 0

INTRODUCTION

NAFLD is emerging as most common liver disorder in India and other developing countries, The histological spectrum ranging from simple steatosis or steatosis with only mild inflammation(type 1 and 2 NAFLD) to more severe steatohepatitis (types 3 and 4 NAFLD or NASH. **Types 3 and 4 NAFLD** progress to cirrhosis in **15-20%** of patients. Progression is silent or paradoxically associated with normalization of aminotransferases. **"NASH"** coined by Ludwig and colleagues from Mayo clinic.

SPECTRUM OF NAFLD

Steatosis

PAGE: 1 OF 70

startDownloadsIMG_0006 ~...Transcend (...KINGSTON (...Turnitin - Go...Turnitin Doc...Task Managerturnitin [Co...mail turnitin ...3:27 PM